Liebenberg syndrome

Liebenberg syndrome is a condition that involves abnormal development of the arms, resulting in characteristic arm malformations that can vary in severity. In people with this condition, bones and other tissues in the elbows, forearms, wrists, and hands have characteristics of related structures in the lower limbs. For example, bones in the elbows are abnormally shaped, which affects mobility of the joints. The stiff elbows function more like knees, unable to rotate as freely as elbows normally do. Bones in the wrists are joined together (fused), forming structures that resemble those in the ankles and heels and causing permanent bending of the hand toward the thumb (radial deviation). The bones in the hands (metacarpals) are longer than normal, and the fingers are short (brachydactyly), similar to the proportions of bones found in the feet. In addition, muscles and tendons that are typically found only in the hands and not in the feet are missing in people with Liebenberg syndrome. Affected individuals also have joint deformities (contractures) that limit movement of the elbows, wrists, and hands. Development of the lower limbs is normal in people with this condition.

Individuals with Liebenberg syndrome have no other health problems related to this condition, and life expectancy is normal.

Frequency

Liebenberg syndrome is a rare condition. Fewer than 10 affected families have been described in the medical literature.

Genetic Changes

Liebenberg syndrome is caused by genetic changes near the *PITX1* gene. The protein produced from this gene plays a critical role in lower limb development by controlling the activity of other genes involved in limb development, directing the shape and structure of bones and other tissues in the legs and feet.

The genetic changes involved in Liebenberg syndrome delete, insert, or rearrange genetic material near the *PITX1* gene. These changes affect regions of DNA known as regulatory elements, which help turn on or turn off genes (known as enhancers or repressors, respectively). They control when and where certain genes are active. The mutations that cause Liebenberg syndrome are thought to relocate enhancers that normally promote the activity of genes involved in upper limb development to be near the *PITX1* gene, where they can promote its activity. Alternatively, the mutations may remove repressors that normally turn off the *PITX1* gene during upper limb development. As a result, the *PITX1* gene is abnormally active during development of the upper limbs. Because the PITX1 protein normally directs lower limb structure,

bones, muscles, and tendons in the arms and hands develop more like those in the legs and feet, leading to the features of Liebenberg syndrome.

Inheritance Pattern

Liebenberg syndrome is inherited in an autosomal dominant pattern, which means having a genetic change that affects the *PITX1* gene on one copy of the chromosome in each cell is sufficient to cause the disorder. In most cases, an affected person has one parent with the condition.

Other Names for This Condition

- brachydactyly-elbow wrist dysplasia syndrome
- brachydactyly with joint dysplasia
- carpal synostosis with dysplastic elbow joints and brachydactyly

Diagnosis & Management

Genetic Testing

 Genetic Testing Registry: Liebenberg syndrome https://www.ncbi.nlm.nih.gov/gtr/conditions/C1861313/

General Information from MedlinePlus

- Diagnostic Tests
 https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

- Encyclopedia: Contracture Deformity https://medlineplus.gov/ency/article/003185.htm
- Encyclopedia: Skeletal Limb Abnormalities https://medlineplus.gov/ency/article/003170.htm
- Health Topic: Arm Injuries and Disorders https://medlineplus.gov/arminjuriesanddisorders.html

Genetic and Rare Diseases Information Center

Brachydactyly elbow wrist dysplasia
 https://rarediseases.info.nih.gov/diseases/966/brachydactyly-elbow-wrist-dysplasia

Educational Resources

- American Society for Surgery of the Hand: Congenital Hand Differences http://www.assh.org/handcare/hand-arm-conditions/Congenital-Differences
- Disease InfoSearch: Brachydactyly Elbow Wrist Dysplasia
 http://www.diseaseinfosearch.org/Brachydactyly+Elbow+Wrist+Dysplasia/920
- MalaCards: liebenberg syndrome http://www.malacards.org/card/liebenberg_syndrome
- Orphanet: Brachydactyly-elbow wrist dysplasia syndrome http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=1275
- The Hospital for Sick Children: Congenital Hand Anomaly (Hand Difference) http://www.aboutkidshealth.ca/En/HealthAZ/ConditionsandDiseases/ MuscleBoneandJointDisorders/Pages/Congenital-Hand-Anomaly.aspx

Patient Support and Advocacy Resources

- Birth Defect Research for Children http://www.birthdefects.org/
- March of Dimes http://www.marchofdimes.org/complications/birth-defects-and-healthconditions.aspx
- Resource List from the University of Kansas Medical Center: Limb Anomalies http://www.kumc.edu/gec/support/limb.html

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28Liebenberg+syndrome %5BTIAB%5D%29+OR+%28SYNOSTOSIS,+CARPAL,+WITH+DYSPLASTIC +ELBOW+JOINTS+AND+BRACHYDACTYLY%5BTIAB%5D%29+OR+ %28BRACHYDACTYLY+WITH+JOINT+DYSPLASIA%5BTIAB%5D%29%29+AND +english%5Bla%5D+AND+human%5Bmh%5D

OMIM

 LIEBENBERG SYNDROME http://omim.org/entry/186550

Sources for This Summary

- Al-Qattan MM, Al-Thunayan A, Alabdulkareem I, Al Balwi M. Liebenberg syndrome is caused by a
 deletion upstream to the PITX1 gene resulting in transformation of the upper limbs to reflect lower
 limb characteristics. Gene. 2013 Jul 15;524(1):65-71. doi: 10.1016/j.gene.2013.03.120. Epub 2013
 Apr 12.
 - Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23587911
- DeLaurier A, Schweitzer R, Logan M. Pitx1 determines the morphology of muscle, tendon, and bones of the hindlimb. Dev Biol. 2006 Nov 1;299(1):22-34. Epub 2006 Jul 14. *Citation on PubMed:* https://www.ncbi.nlm.nih.gov/pubmed/16989801
- Duboc V, Logan MP. Pitx1 is necessary for normal initiation of hindlimb outgrowth through regulation of Tbx4 expression and shapes hindlimb morphologies via targeted growth control. Development. 2011 Dec;138(24):5301-9. doi: 10.1242/dev.074153. Epub 2011 Nov 9. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/22071103
 Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3222209/
- Seoighe DM, Gadancheva V, Regan R, McDaid J, Brenner C, Ennis S, Betts DR, Eadie PA, Lynch SA. A chromosomal 5q31.1 gain involving PITX1 causes Liebenberg syndrome. Am J Med Genet A. 2014 Nov;164A(11):2958-60. doi: 10.1002/ajmg.a.36712. Epub 2014 Aug 14. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/25124102
- Spielmann M, Brancati F, Krawitz PM, Robinson PN, Ibrahim DM, Franke M, Hecht J, Lohan S, Dathe K, Nardone AM, Ferrari P, Landi A, Wittler L, Timmermann B, Chan D, Mennen U, Klopocki E, Mundlos S. Homeotic arm-to-leg transformation associated with genomic rearrangements at the PITX1 locus. Am J Hum Genet. 2012 Oct 5;91(4):629-35. doi: 10.1016/j.ajhg.2012.08.014. Epub 2012 Sep 27.

Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/23022097
Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3484647/

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/liebenberg-syndrome

Reviewed: September 2016 Published: March 21, 2017 Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services